

Large Scale Traveling Ionospheric Disturbances Observed using HamSCI Amateur Radio, SuperDARN, GNSS TEC, and Ionosondes.

Diego F. Sanchez KD2RLM^{1,3}, Nathaniel A. Frissell W2NAF², Gareth W. Perry KD2SAK³, William D. Engelke AB4EJ⁴, Anthea Coster⁵, Philip J. Erickson W1PJE⁵, J. Michael Ruohoniemi⁶, and Joseph Baker⁶, Carl Luetzelschwab K9LA

¹Essex County College

²University of Scranton

³New Jersey Institute of Technology

⁴University of Alabama

⁵MIT Haystack Observatory

⁶Virginia Tech


Traveling Ionospheric Disturbances

- TIDs are Quasi-periodic Variations of F Region Electron Density
- Medium Scale (MSTID)
 - $T \approx 15 - 60$ min
 - $v_H \approx 100 - 250$ m/s
 - $\lambda_H \approx$ Several Hundred km (< 1000 km)
 - Often Meteorological Sources
- Large Scale (LSTID)
 - $\lambda_h > 1000$ km
 - $30 < T [\text{min}] < 180$
 - Often Auroral Electrojet Enhancement, Particle Precipitation
- Often associated with Atmospheric Gravity Waves

[Francis, 1975; Hunsucker 1982; Ogawa et al., 1967; Ding et al., 2012; Frissell et al., 2014;
2016]


Traveling Ionospheric Disturbances

MSTIDs are a type of HF Fading



[Samson et al., 1990]

Traveling Ionospheric Disturbances



Ray trace simulation illustrating how SuperDARN HF radars observe MSTIDs.



- (a) Fort Hays East (FHE) radar field of view superimposed on a 250 km altitude cut of a perturbed IRI. FHE Beam 7 is outlined in bold.
- (b) Vertical profile of 14.5 MHz ray trace along FHE Beam 7. Background colors represent perturbed IRI electron densities. The areas where rays reach the ground are potential sources of backscatter.
- (c) Simulated FHE Beam 7 radar data, color coded by radar backscatter power strength. Periodic, slanted traces with negative slopes are the signatures of MSTIDs moving toward the radar.

[\[Frissell et al., 2016\]](#)

Data Sources



- **Ham Radio**
 - Reverse Beacon Network
 - Weak Signal Propagation Reporting Network (WSPRNet)
 - QRZ.com
- **SuperDARN**
- **Madrigal GPS Total Electron Content (TEC)**
- **NASA OMNI Data**
- **Ionosonde foF2 Data**

TIDs in Blackstone SuperDARN




Ham Radio TIDs


N Spots = 157559
RBN: 29%
WSPRNet: 71%



GNSS TEC Comparison 18:00 - 21:00






- Radio range is shortest when TEC is red (higher TEC)
- Higher electron densities → More HF refraction, communication range decreases






November 3, 2017 US Ham Radio TID

23 Nov 2017-
24 Nov 2017
Ham Radio Networks
N Spots = 175674
RBN: 38%
WSPRNet: 62%






Nov 23, 2017 GNSS TEC Comparison

23 Nov 2017-
24 Nov 2017
Ham Radio Networks
N Spots = 175674
RBN: 38%
WSPRNet: 62%






Nov 23, 2017 GNSS TEC Comparison

23 Nov 2017-
24 Nov 2017
Ham Radio Networks
N Spots = 175674
RBN: 38%
WSPRNet: 62%






Nov 23, 2017 GNSS TEC Comparison

23 Nov 2017-
24 Nov 2017
Ham Radio Networks
N Spots = 175674
RBN: 38%
WSPRNet: 62%






Nov 23, 2017 GNSS TEC Comparison

23 Nov 2017-
24 Nov 2017
Ham Radio Networks
N Spots = 175674
RBN: 38%
WSPRNet: 62%






January 26, 2017 US Ham Radio TID

26 Jan 2017-
27 Jan 2017
Ham Radio Networks
N Spots = 151871
RBN: 48%
WSPRNet: 52%







Jan 26, 2017 GNSS TEC Comparison

26 Jan 2017-
27 Jan 2017
Ham Radio Networks
N Spots = 151871
RBN: 48%
WSPRNet: 52%







Jan 26, 2017 GNSS TEC Comparison

26 Jan 2017-
27 Jan 2017
Ham Radio Networks
N Spots = 151871
RBN: 48%
WSPRNet: 52%







Jan 26, 2017 GNSS TEC Comparison

26 Jan 2017-
27 Jan 2017
Ham Radio Networks
N Spots = 151871
RBN: 48%
WSPRNet: 52%







Jan 26, 2017 GNSS TEC Comparison

26 Jan 2017-
27 Jan 2017
Ham Radio Networks
N Spots = 151871
RBN: 48%
WSPRNet: 52%




Jan 26, 2017 GNSS TEC Comparison

26 Jan 2017-
27 Jan 2017
Ham Radio Networks
N Spots = 151871
RBN: 48%
WSPRNet: 52%



Ionosonde Data





Juliusruh Ionosonde
Rügen, Germany

- Chose ionosondes relatively close to the largest concentration of radio spots.





Boulder Ionosonde
Colorado, USA



January 26, 2017 foF2 Comparison US



14 MHz



- The highest frequency that the ionosonde reflects is called foF2.
- The higher the frequency that is bounced back from the ionosphere, the higher the electron densities will be.
- More refraction at higher electron densities.
- Radio range is shortest at higher foF2 values.

September 27, 2017 foF2 Comparison Europe

14 MHz



- The highest frequency that the ionosonde reflects is called foF2.
- The higher the frequency that is bounced back from the ionosphere, the higher the electron densities will be.
- More refraction at higher electron densities.
- Radio range is shortest at higher foF2 values.

Conclusions and Future Work

- **Ham Radio Disturbances Seem likely to be LSTIDs**
 - Appear coherent across Continental US and Europe
 - Consistent with BKS SuperDARN Beam 13
 - Consistent with GNSS TEC
 - Consistent with Boulder (US) and Juliusruh (Europe) Ionosondes.
- **RBN and WSPR can serve as a tool for monitoring LSTIDs day and night**
- **Find exact source mechanism of the detected LSTIDs.**

References

Ding, F., Wan, W., Ning, B., Zhao, B., Li, Q., Zhang, R., Xiong, B., and Song, Q. (2012), Two-dimensional imaging of large-scale traveling ionospheric disturbances over China based on GPS data, *J. Geophys. Res.*, 117, A08318, doi:10.1029/2012JA017546.

Francis, S. H. (1975), Global propagation of atmospheric gravity waves: A review, *J. Atmos. Terr. Phys.*, 37, 1011–1054, doi:10.1016/0021-9169(75)90012-4.

Frissell, N. A., Baker, J. B. H., Ruohoniemi, J. M., Gerrard, A. J., Miller, E. S., Marini, J. P., West, M. L., and Bristow, W. A. (2014), Climatology of medium-scale traveling ionospheric disturbances observed by the midlatitude Blackstone SuperDARN radar, *J. Geophys. Res. Space Physics*, 119, 7679– 7697, doi:10.1002/2014JA019870.

Frissell, N. A., Baker, J. B. H., Ruohoniemi, J. M., Greenwald, R. A., Gerrard, A. J., Miller, E. S., and West, M. L. (2016), Sources and characteristics of medium-scale traveling ionospheric disturbances observed by high-frequency radars in the North American sector, *J. Geophys. Res. Space Physics*, 121, 3722– 3739, doi:10.1002/2015JA022168.





Chimonas, G. (1970), The equatorial electrojet as a source of long period traveling ionospheric disturbances, *Planet. Space Sci.*, 18(4), 583–589, doi:10.1016/0032-0633(70)90133-9.

Vadas, S. L., and H. Liu (2009), Generation of large-scale gravity waves and neutral winds in the thermosphere from the dissipation of convectively generated gravity waves, *J. Geophys. Res.*, 114, A10310, doi:10.1029/2009JA014108.

Extra Slides




01/26/2017 - 01/27/2017 European Response

26 Jan 2017-
27 Jan 2017
Ham Radio Networks
N Spots = 321853
RBN: 30%
WSPRNet: 70%








05/02/2017 - 05/03/2017 US Response

02 May 2017-
03 May 2017
Ham Radio Networks
N Spots = 161859
RBN: 17%
WSPRNet: 83%







05/02/2017 - 05/03/2017 European Response

02 May 2017-
03 May 2017
Ham Radio Networks
N Spots = 381508
RBN: 16%
WSPRNet: 84%








09/27/2017 - 09/28/2017 US Response

27 Sep 2017-
28 Sep 2017
Ham Radio Networks
N Spots = 192946
RBN: 12%
WSPRNet: 88%








09/30/2017 - 10/01/2017 US Response

30 Sep 2017-
01 Oct 2017
Ham Radio Networks
N Spots = 170377
RBN: 11%
WSPRNet: 89%








09/30/2017 - 10/01/2017 European Response

30 Sep 2017-
01 Oct 2017
Ham Radio Networks
N Spots = 527440
RBN: 24%
WSPRNet: 76%




11/23/2017 - 11/24/2017 European Response

**23 Nov 2017-
24 Nov 2017**
Ham Radio Networks
N Spots = 284419
RBN: 21%
WSPRNet: 79%




01/26/2017 - 01/27/2017 US Response


26 Jan 2017-
27 Jan 2017
Ham Radio Networks
N Spots = 151871
RBN: 48%
WSPRNet: 52%



Radio Spots (N = 57548)



Radio Spots (N = 94323)



09/27/2017 - 09/28/2017 European Response

